Oy Medix Biochemica Ab
Headquarters
Klovinpellontie 3
FI-02180 Espoo
Finland
Email: sales@medixbiochemica.com

Medix Biochemica USA Inc.
10850 Metro Court
Maryland Heights, MO 63043
United States of America
Email: sales@medixbiochemica.com

Product Datasheet

Human IL-13 ELISA Kit
850.080

Oy Medix Biochemica Ab
Headquarters
Klovinpellontie 3
FI-02180 Espoo
Finland
Email: sales@medixbiochemica.com

Medix Biochemica USA Inc.
10850 Metro Court
Maryland Heights, MO 63043
United States of America
Email: sales@medixbiochemica.com

Product Name	Human IL-13 ELISA Kit
Catalog Number	850.080
Description	Diaclone ELISA Kits include pre-coated strip plates, biotinylated secondary antibody, standards, controls (where applicable), buffers, streptavidin-HRP, TMB, stop reagents, and a detailed procedure.
Range	3.125-100 pg/ml
Sensitivity	1.5 pg/ml
Brand	Diaclone
Storage	Store at +4°C
Cross Reactivity	No cross reactivity with other human cytokines
Specificity	Recognizes both natural and recombinant human IL-13
Sample Types	Serum, Plasma, Cell culture supernatant

Oy Medix Biochemica Ab
Headquarters
Klovinpellontie 3
FI-02180 Espoo
Finland
Email: sales@medixbiochemica.com

Medix Biochemica USA Inc.
10850 Metro Court
Maryland Heights, MO 63043
United States of America
Email: sales@medixbiochemica.com

References

Piccoli L. et al. Clin Dev Immunol. 2012: 483935 - **Pubmed link**
Malhotra, N. et al., Lung India. 2020 Jul-Aug; 37(4): 310–315. - **Pubmed link**
Sarbinowska, J. et al., Biomolecules. 2021 Jun; 11(6): 890. - **Pubmed link**
Qurashi, T.A. et al., Allergy Asthma Clin Immunol. 2021; 17(1): 119. - **Pubmed link**
Abbal C. et al., Int. Immunol., 1999; 11(9) : 1451 - 1462 - **Pubmed link**
Schmitt, C. et al., J Leukoc Biol.,2000; 68(6): 836-44. - **Pubmed link**
Saverino D. et al., J. immunol., 2002; 168(1): 207 - 215 - **Pubmed link**
Vekemans, J. et al.,Clin Exp Immunol.,2002;127(3):495-8 - **Pubmed link**
Fahy O. et al., J. immunol., 2002; 168(11) : 5912 - 5919 - **Pubmed link**
Woerly, G. et al., J Leukoc Biol.,2002; 72(4): 769-79. - **Pubmed link**
Ismaili, J. et al., Clin Exp Immunol.,2003;133(3):414-21 - **Pubmed link**
Leung S. et al., J. Biomol Screen, 2005; 10(2):157 - 167 - **Pubmed link**
de Nadai, P. et al., J Immunol.,2006;176(10): 6286-93. - **Pubmed link**
Cameron, L. et al., J Immunol.,2006;177(12): 8633-42. - **Pubmed link**
Gafvelin, G. et al., J Biol Chem., 2007; 282(6): 3778-87. - **Pubmed link**
Pene, J. et al., J. Immunol.,2008; 180(11): 7423-7430. - **Pubmed link**
Finan, C. et al., PLoS One,2008; 3(10): e3485 - **Pubmed link**
Halonen, M. et al., J. Immunol.,2009;182(5): 3285-3293. - **Pubmed link**
Pitta, M. G. et al., J Clin Invest.,2009; 119(8): 2379-87 - **Pubmed link**
Perrin, Y. et al.,Clin Transl Allergy,2014; 4(1): 1 - **Pubmed link**
Tsvetkova-Vicheva, V. M. et al.,Clin Transl Allergy,2014; 4(1): 3 - **Pubmed link**
Antczak A. et al. International Journal of Immunopathology and Pharmacology 2016: 0394632015623794 - **Pubmed link**
Knuhr, K. et al., Front Immunol. 2018 Oct 10;9:2293 - **Pubmed link**