



Product specifications

Name	Anti-h CA125 4602 SPTN-5			
Specificity	Antibody recognizes human cancer antigen 125 (CA125)			
Description	Monoclonal mouse antibody, cultured <i>in vitro</i> under conditions free from animal-derived components.			
Product code	100628			
Product buffer solution	50 mM Na-citrate, pH 6.0, 0.9 % NaCl, 0.095 % NaN $_3$ as a preservative			
Shelf life and storage	12 months from manufacturing at 2–8 °C			
Subclass	IgG1			
Analyte description	Cancer antigen 125 is a peptide epitope on the mucin glycoprotein MUC16. CA125 is the most widely used serum biomarker to monitor epithelial ovarian cancer. It is also used for the differential diagnosis of pelvic masses.			

Parameters tested on each lot

Product appearance	Liquid, may turn slightly opaque during storage
Product concentration	5.0 mg/ml (+/- 10 %)
Immunoreactivity	80–120 % compared to the reference sample in an FIA test
IEF Profile	6.3–7.3
Purity	≥ 95 %

Kinetic parameters

Association rate constant	Not Determined (N/D)
Dissociation rate constant	N/D
Affinity constant	N/D
Determination method	-
Determination antigen	

Oy Medix Biochemica Ab Klovinpellontie 3 FI-02180 Espoo, Finland

medix@medixbiochemica.com www.medixbiochemica.com VAT reg.no. FI14631532

Legal disclaimer Medix Biochemica products meet their specifications if transported, stored and used according to the instructions. Medix/MAB and Medix/Antigens are trademarks of Medix Biochemica and may not be used or reproduced without Medix Biochemica's written permission.





Cross-reactivities	Does not recognize carbohydrate antigen 19-9 (CA19-9), cancer antigen 15-3 (CA15-3), CYFRA 21-1, carcinoembryonic antigen (CEA), α-fetoprotein (AFP), or human epididymis protein 4 (HE4).					
Epitope	The antibody 4602 is M11-like and belongs to group B (ISOBM classification) as described in Nustad et al. 1996.					
Pair recommendations	DETECTION		TION			
		4601	4602			
	Harrison 4601 Harrison 4602	-	+			
	4602	+	-			
	Please note that pair recommendations are based on results obtained by our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations are only indicative.					
Platforms tested	FIA					
Antigens tested	Native CA125 antigen, Lee Biosolutions 151-25.					
Product stability	TEMPERATURE, TIME			RESULT		
	-70 °C, 21 days			OK		
	-20 °C, 21 days			OK		
	+4 °C, 21 days			ОК		
	+35 °C, 21 days			OK		
	+45 °C, 7 days			ОК		
	Stability testing is performed in the product buffer to see whether different temperatures affect the antigen binding, charge or composition of the antibody. Please note that the shelf life given on the first page is based on real time stability testing at 2–8 °C in the product buffer.					
Miscellaneous	-					
References	Nustad K, Bast RC, O'Brien TJ et al. (1996) Specificity and affinity of 26 monoclonal antibodies against the CA 125 antigen: First report from the ISOBM TD-1 Workshop. Tumor Biol 17:196- 219.					

Oy Medix Biochemica Ab Klovinpellontie 3 FI-02180 Espoo, Finland

medix@medixbiochemica.com www.medixbiochemica.com VAT reg.no. FI14631532

Legal disclaimer Medix Biochemica products meet their specifications if transported, stored and used according to the instructions. MedixMAB and MedixAntigens are trademarks of Medix Biochemica and may not be used or reproduced without Medix Biochemica's written permission.