



Product specifications

Name Anti-h CYFRA21-1 1605 SPTN-5

Specificity Antibody recognizes human CYFRA 21-1 fragment of cytokeratin 19

Description Monoclonal mouse antibody, cultured *in vitro* under conditions free from animal-derived

components

Product code 100223

Product buffer solution 50 mM Na-citrate, pH 6.0, 0.9 % NaCl, 0.095 % NaN₃ as a preservative

Shelf life and storage 12 months from manufacturing at 2–8 °C

Subclass IgG₁

Analyte description CYFRA 21-1 is a fragment of cytokeratin 19 which is expressed in various types of epithelial

cells and tumor cells of epithelial origin. Serum CYFRA 21-1 has been used as a tumor marker

of different origins and especially in non-small cell lung cancer.

Parameters tested on each lot

Product appearance Liquid, may turn slightly opaque during storage

Product concentration 5.0 mg/ml (+/- 10 %)

Immunoreactivity 80–120 % compared to the reference sample in an FIA test

IEF Profile 6.7–7.4

Purity ≥ 95 %

Kinetic parameters

Association rate constant 5 x 10⁴ 1/Ms

Dissociation rate constant 6 x 10⁻⁵ 1/s

Affinity constant $K_A = 8 \times 10^8 \text{ 1/M}, K_D = 1 \times 10^{-9} \text{ M}$

Determination method SPR analysis (ProteOn XPR36)

Determination antigen Recombinant human Cytokeratin 19, Biodesign (Cat A44214H, Lot 12F15608)





2021-04-21

Cross-reactivities 100 % cytokeratin 19

Epitope Not Determined (N/D)

Pair recommendations

		DETECTION			
		1602	1603	1604	1605
CAPTURE	1602	-	-	+	+
	1603	-	-	+	+
	1604	+	+	-	-
	1605	+	+	-	-

Please note that pair recommendations are based on results obtained by our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations are only indicative.

Platforms tested FIA

Antigens tested Native CYFRA 21-1 antigen, Lee Biosolutions 192-20

Product stability TEMPERATURE, TIME RESULT

-70 °C, 21 days OK
-20 °C, 21 days OK
+4 °C, 21 days OK
+25 °C, 21 days OK
+35 °C, 21 days OK
+45 °C, 7 days OK

Stability testing is performed in the product buffer to see whether different temperatures affect the antigen binding, charge or composition of the antibody. Please note that the shelf life given on the first page is based on real time stability testing at 2–8 °C in the product buffer.

Miscellaneous -

References -