

## Product specifications

---

Name	Anti-h Prealbumin 11605 SPTN-5
Specificity	Antibody recognizes human prealbumin
Description	Monoclonal mouse antibody, cultured <i>in vitro</i> under conditions free from animal-derived components
Product code	100832
Product buffer solution	50 mM Na-citrate, pH 6.0, 0.9 % NaCl, 0.095 % NaN <sub>3</sub> as a preservative
Shelf life and storage	Unspecified, storage at 2–8 °C
Subclass	IgG <sub>1</sub>
Analyte description	Prealbumin, also known as transthyretin, is synthesized mainly by liver. It is involved in transport of triiodothyronine (T3), thyroxine (T4), and vitamin A. Due to the short half-life of prealbumin, it can be used to assess malnutrition or efficacy of parenteral nutrition.

## Parameters tested on each lot

---

Product appearance	Liquid, may turn slightly opaque during storage
Product concentration	5.0 mg/ml (+/- 10 %)
Immunoreactivity	80–120 % compared to the reference sample in an FIA test
IEF Profile	6.3–6.7
Purity	≥ 95 %

## Kinetic parameters

---

Association rate constant	Not Determined (N/D)
Dissociation rate constant	N/D
Affinity constant	N/D
Determination method	-
Determination antigen	-

### Legal disclaimer



Cross-reactivities N/D

Epitope N/D

Pair recommendations

		DETECTION				
		11601	11603	11604	11605	11606
CAPTURE	11601	+	+	+	+	+
	11603	+	+	+	+	+
	11604	+	+	+	+	+
	11605	+	+	+	+	+
	11606	+	+	+	+	+

Following pairs are especially recommended for the below mentioned assays:  
IT: 11601 – 11604, 11601 – 11605 and 11604 – 11605

Please note that pair recommendations are based on results obtained by our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations are only indicative.

Platforms tested FIA, IT

Antigens tested Native prealbumin antigen Lee Biosolutions 496-10 and 496-11.

TEMPERATURE, TIME	RESULT
-70 °C, 21 days	OK
-20 °C, 21 days	OK
+4 °C, 21 days	OK
+35 °C, 21 days	OK
+45 °C, 7 days	OK

Stability testing is performed in the product buffer to see whether different temperatures affect the antigen binding, charge or composition of the antibody. Please note that the shelf life given on the first page is based on real time stability testing at 2–8 °C in the product buffer.

Miscellaneous -

References -

