



Product specifications

Name Anti-h cTnT 3710 SPTN-5

Specificity Antibody recognizes human cardiac troponin T (cTnT)

Description Monoclonal mouse antibody, cultured *in vitro* under conditions free from animal-derived

components

Product code 100698

Product buffer solution 50 mM Na-citrate, pH 6.0, 0.9 % NaCl, 0.095 % NaN₃ as a preservative

Shelf life and storage Unspecified, storage at 2–8 °C

Subclass IgG₁

Analyte description Troponins form a complex of three regulatory proteins (troponin C, I and T) that are integral

to muscle contraction in skeletal and cardiac muscles. Serum cardiac troponin tests can be used to help diagnose several different heart disorders, especially myocardial infarction.

Parameters tested on each lot

Product appearance Liquid, may turn slightly opaque during storage

Product concentration 5.0 mg/ml (+/- 10 %)

Immunoreactivity 80–120 % compared to the reference sample in an FIA test

IEF Profile 6.5–7.0

Purity ≥ 95 %

Kinetic parameters

Association rate constant Not Determined (N/D)

Dissociation rate constant N/D

Affinity constant N/D

Determination method -

Determination antigen -





2021-04-21

Cross-reactivities

Does not recognize skeletal troponin T (sTnT) or cardiac troponin I (cTnI).

Epitope

Amino acids 139-152 (RIRNEREKERQNRL) of cardiac troponin T isoform 6.

Pair recommendations

		DETECTION					
		3701	3703	3708	3710	3711	3712
CAPTURE	3701	-	-	ı	-	-	-
	3703	-	-	-	+	+	+
	3708	+	+	-	+	+	+
	3710	-	+	+	-	-	-
	3711	-	+	-	-	-	-
	3712	-	+	+	-	-	-

Please note that pair recommendations are based on results obtained by our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations are only indicative.

Platforms tested

FIA

Antigens tested

Recombinant cardiac troponin T antigen, Medix Biochemica 610101 and native cardiac troponin T antigen, Lee Biosolutions 550-21.

Product stability

TEMPERATURE, TIME	RESULT
-70 °C, 21 days	OK
-20 °C, 21 days	OK
+4 °C, 21 days	OK
+35 °C, 21 days	OK
+45 °C, 7 days	OK

Stability testing is performed in the product buffer to see whether different temperatures affect the antigen binding, charge or composition of the antibody. Please note that the shelf life given on the first page is based on real time stability testing at 2–8 °C in the product buffer.

Miscellaneous

References