

Product specification ANTIBODY

2022-11-02

Anti-h Cystatin C 10002 SPTN-5

Product overview

Catalog number 100690

Specificity Antibody recognizes human Cystatin C

Description Monoclonal mouse antibody, cultured *in vitro* under conditions free from

animal-derived components.

Product buffer solution 50 mM Na-citrate, pH 6.0, 0.9 % NaCl, 0.095 % NaN₃ as a preservative

Shelf life and storage Unspecified, storage at 2–8 °C

 $\label{eq:G2b} \textbf{Subclass} \qquad \qquad \textbf{IgG}_{2b}$

Analyte description Cystatin C is an emerging renal biomarker. It is used for the diagnosis of

chronic kidney disease. Cystatin C has also been associated with an

increased risk of cardiovascular disease and heart failure.

Parameters tested on each lot

Product appearance Liquid, may turn slightly opaque during storage

Product concentration 5.0 mg/ml (+/- 10 %)

Immunoreactivity 80–120 % compared to the reference sample in an FIA test

IEF Profile 6.2–7.3

Purity $\geq 95\%$

Kinetic parameters

Association rate constant Not Determined (N/D)

Dissociation rate constant N/D

Affinity constant N/D

Determination method -

Determination antigen -





Product specification ANTIBODY

2022-11-02

Cross-reactivities Does not recognize recombinant Cystatin D, F, S, SA, or SN.

Epitope N/D

Pair recommendations

		DETECTION			
		10001	10002	10004	10005
CAPTURE	10001	-	-	+	+
	10002	-	-	+	+
	10004	+	+	-	-
	10005	+	+	-	-

Please note that pair recommendations are based on results obtained by our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations are only indicative.

Platforms tested FIA

Antigens tested Recombinant Cystatin C antigen, Medix Biochemica, 610100.

Product stability TEMPERATURE, TIME RESULT

-70 °C, 21 days OK
-20 °C, 21 days OK
+4 °C, 21 days OK
+35 °C, 21 days OK
+45 °C, 7 days OK

Stability testing is performed in the product buffer to see whether different temperatures affect the antigen binding, charge or composition of the antibody. Please note that the shelf life given on the first page is based on real time stability testing at 2–8 °C in the product buffer.

Miscellaneous -

References -

