

## Anti-h IgE 8518 SPRNE-1

### Product overview

---

<b>Catalog number</b>	100106
<b>Specificity</b>	Antibody recognizes human immunoglobulin E
<b>Description</b>	Monoclonal mouse antibody, cultured <i>in vitro</i> under conditions free from animal-derived components.
<b>Product buffer solution</b>	37 mM citrate, 125 mM phosphate, pH 6.0, 0.9 % NaCl, 25 % ethylene glycol, 0.095 % NaN <sub>3</sub> as a preservative
<b>Shelf life and storage</b>	36 months from manufacturing at 2–8 °C
<b>Subclass</b>	IgG <sub>1</sub>
<b>Analyte description</b>	Immunoglobulin E (IgE) is a class of antibody that has only been found in mammals. It plays an important role in allergy and defence against parasites. Although IgE is typically the least abundant isotype - blood serum IgE levels in a normal individual are only 0.05 % of the IgG concentration - it is capable of triggering the most powerful immune reactions.

### Parameters tested on each lot

---

<b>Product appearance</b>	Liquid, may turn slightly opaque during storage
<b>Product concentration</b>	1.0 mg/ml (+/- 10 %)
<b>Immunoreactivity</b>	80–120 % compared to the reference sample in an FIA test
<b>IEF Profile</b>	6.3–7.8
<b>Purity</b>	≥ 95 %

### Kinetic parameters

---

<b>Association rate constant</b>	Not Determined (N/D)
<b>Dissociation rate constant</b>	N/D
<b>Affinity constant</b>	$K_A = 1.2 \times 10^{10} \text{ 1/M}$
<b>Determination method</b>	SPR analysis (Biacore)
<b>Determination antigen</b>	IgE, Biodesign (Cat A101644, Lot 912737)



#### Legal disclaimer

**Cross-reactivities** Human IgA <0.08 %  
Human IgG <0.08 %  
Human IgM <0.08 %

**Epitope** N/D

**Pair recommendations**

		DETECTION		
		8510	8516	8518
CAPTURE	8510	-	+	+
	8516	-	-	-
	8518	-	-	-

Please note that pair recommendations are based on results obtained by our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations are only indicative.

**Platforms tested** FIA

**Antigens tested** N/D

Product stability	TEMPERATURE, TIME	RESULT
	-70 °C, 21 days	N/D
	-20 °C, 21 days	N/D
	+4 °C, 21 days	N/D
	+30 °C, 7 days	N/D
	+35 °C, 21 days	N/D
	+45 °C, 7 days	N/D

Stability testing is performed in the product buffer to see whether different temperatures affect the antigen binding, charge or composition of the antibody. Please note that the shelf life given on the first page is based on real time stability testing at 2–8 °C in the product buffer.

**Miscellaneous** -

**References** -



**Legal disclaimer**