Medix Biochemica

Anti-h TSH 5409 SPTNE-5

Product overview							
Catalog number	100034						
Specificity	Antibody recognizes human thyrotropin						
Description	Monoclonal mouse antibody, cultured <i>in vitro</i> under conditions free fron animal-derived components.						
Product buffer solution	50 mM Na-citrate, pH 6.0, 0.9 % NaCI, 25 % ethylene glycol, 0.095 % NaN_3 as a preservative						
Shelf life and storage	36 months from manufacturing at 2–8 °C						
Subclass	lgG₁						
Analyte description	Thyroid-stimulating hormone (also known as TSH or thyrotropin) is a peptide hormone synthesized and secreted by thyrotrope cells in the anterior pituitary gland which regulates the endocrine function of the thyroid gland. TSH levels are tested in the blood of patients suspected of suffering from excess (hyperthyroidism), or deficiency (hypothyroidism) o thyroid hormone.						
Parameters tested on each lo	t						
Product appearance	Liquid, may turn slightly opaque during storage						
Product concentration	5.0 mg/ml (+/- 10 %)						
Immunoreactivity	80–120 % compared to the reference sample in an FIA test						
IEF Profile	5.8–7.2						
Purity	≥ 95 %						
Kinetic parameters							
Association rate constant	2.1 x 10 ⁶ 1/Ms						
Dissociation rate constant	1.0 x 10 ⁻⁴ 1/s						
Affinity constant	$K_A = 2.1 \text{ x } 10^{10} \text{ 1/M}; K_D = 4.8 \text{ x } 10^{-11} \text{ M} (= 0.05 \text{ nM})$						
Determination method	SPR analysis (ProteOn XPR36)						
Determination antigen	TSH, Scripps (Cat T0114, Lot 2414402)						



Oy Medix Biochemica Ab Klovinpellontie 3, Fl-02180 Espoo, Finland medix@medixbiochemica.com www.medixbiochemica.com VAT reg. no. FI14631532

Legal disclaimer

Medix Biochemica products meet their specifications if transported, stored, and used according to the instructions. Medix Biochemica's products may not be used or reproduced without Medix Biochemica's written permission.

Medix Biochemica

Cross-reactivities	TSH α TSH β hCG	 < 0.01 % (Scripps Laboratories, Cat T0214, Lot BE298001) 0.14 % (Scripps Laboratories, Cat T0314, Lot BA317002) < 0.04 % (Scripps Laboratories, Cat C0714, Lot BB274002) < 0.04 % (Scripps Laboratories, Cat L0815, Lot BB274002)
	LH	< 0.04 % (Scripps Laboratories, Cat L0815, Lot BB236003)
	FSH	< 0.04 % (Scripps Laboratories, Cat F0614, Lot BA127003)

Epitope

Group 2. The epitope is located at the junction of alpha and beta subunits in hTSH. The antibody recognizes intact hTSH molecule.

Ρ

Pair recommendations			DETECTION]		
			5401	5404	5405	5407	5408	5409			
		5401	-	-	+	+	-	+			
	ш	5404	-	-	+	+	-	+			
	CAPTURE	5405	-	-	-	-	+	+			
	AP	5407	-	-	-	-	+	+	_		
		5408	-	-	+	+	-	+	_		
		5409	-	-	+	+	-	-			
	 Following pairs are especially recommended for the below mentioned assays: CLIA: 5405 (capture) – 5409 (detection), 5407 – 5409 and 5409 – 540' LF: 5405 (membrane) – 5409 (particles), 5407 – 5409, 5408 – 5405, 5408 – 5407, 5408 – 5409, 5409 – 5407 Please note that pair recommendations are based on results obtained our laboratory. Equally good results may be obtained using other pairs therefore these recommendations are only indicative. 										
Platforms tested	FIA, LF, CLIA										
Antigens tested	Native TSH antigen Lee Biosolutions 996-50 and 996-51.										
Product stability	-20 °C, 14 daysO+4 °C, 14 daysO+35 °C, 14 daysO+45 °C, 7 daysOStability testing is performed in the proc temperatures affect the antigen binding antibody. Please note that the shelf life						ermined uffer to so ge or cor on the fil	ee wheth nposition rst page	of the		
Miscellaneous	real time stability testing at 2–8 °C in the product buffer.										
References	Brockmann, EC., Pettersson, K and Vehniäinen M. (2010) Use of high- capacity surface with oriented recombinant antibody fragments in a 5-min immunoassay for thyroid-stimulating hormone. Anal. Biochem., 396:242- 249										



Oy Medix Biochemica Ab Klovinpellontie 3, FI-02180 Espoo, Finland

medix@medixbiochemica.com www.medixbiochemica.com VAT reg. no. FI14631532

Legal disclaimer

Medix Biochemica products meet their specifications if transported, stored, and used according to the instructions. Medix Biochemica's products may not be used or reproduced without Medix Biochemica's written permission.