# Medix Biochemica

### Anti-h Inhibin alpha 9401 SPTN-5

Product overview		
Catalog number	100711	
Specificity	Antibody recognizes human inhibin alpha subunit	
Description	Monoclonal mouse antibody, cultured <i>in vitro</i> under conditions free from animal-derived components.	
Product buffer solution	50 mM Na-citrate, pH 6.0, 0.9 % NaCl, 0.095 % NaN $_{\rm 3}$ as a preservative	
Shelf life and storage	12 months from manufacturing at 2–8 °C	
Subclass	lgG <sub>1</sub>	
Analyte description	Inhibin alpha can form dimers either with inhibin beta A or inhibin beta B. These dimers are called inhibin A, or inhibin B, respectively. Both inhibin A and inhibin B inhibit FSH secretion. Inhibin A tests can be used in the diagnostics of Trisomy 21 and Trisomy 18. Inhibin B is a marker of male infertility.	
Parameters tested on each lot		

Product appearance	Liquid, may turn slightly opaque during storage	
Product concentration	5.0 mg/ml (+/- 10 %)	
Immunoreactivity	80–120 % compared to the reference sample in an FIA test	
IEF Profile	6.3–7.2	
Purity	≥ 95 %	
Kinetic parameters		
Association rate constant	Not Determined (N/D)	
Dissociation rate constant	N/D	
Affinity constant	N/D	
Determination method	-	
Determination antigen	-	



**Oy Medix Biochemica Ab** Klovinpellontie 3, Fl-02180 Espoo, Finland medix@medixbiochemica.com www.medixbiochemica.com VAT reg. no. FI14631532

### Legal disclaimer

Medix Biochemica products meet their specifications if transported, stored, and used according to the instructions. Medix Biochemica's products may not be used or reproduced without Medix Biochemica's written permission.

# Medix Biochemica

N/D

Epitope

Amino acids 233-264 (STPLMSWPWSPSALRLLQRPPEEPAAHANCHR)

**Cross-reactivities** 

		DETECTION	
		9401	9504
CAPTURE	9401	-	+
	9504	+	-

Please note that pair recommendations are based on results obtained by our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations are only indicative.

Platforms tested	FIA		
Antigens tested	N/D		
Product stability	TEMPERATURE, TIME -70 °C, 21 days -20 °C, 21 days +4 °C, 21 days +35 °C, 21 days +45 °C, 7 days	RESULT OK OK OK OK	
	Stability testing is performed in the product buffer to see whether temperatures affect the antigen binding, charge or composition of antibody. Please note that the shelf life given on the first page is I real time stability testing at 2–8 °C in the product buffer.		

Miscellaneous	
References	



**Oy Medix Biochemica Ab** Klovinpellontie 3, Fl-02180 Espoo, Finland medix@medixbiochemica.com www.medixbiochemica.com VAT reg. no. FI14631532

#### Legal disclaimer

Medix Biochemica products meet their specifications if transported, stored, and used according to the instructions. Medix Biochemica's products may not be used or reproduced without Medix Biochemica's written permission.