

PRODUCT SPECIFICATIONS

Name	Anti-h C-Peptide 9103 SPRN-5
Specificity	Antibody recognizes human C-peptide
Description	Monoclonal mouse antibody, cultured <i>in vitro</i> under conditions free from animal-derived components
Product code	100114
Product buffer solution	37 mM citrate, 125 mM phosphate, pH 6.0, 0.9 % NaCl, 0.095 % NaN ₃ as a preservative
Shelf life and storage	24 months from manufacturing at 2-8 °C
Analyte description	C-peptide is produced when proinsulin is split into insulin and C-peptide. They split before proinsulin is released from endocytic vesicles within the pancreas—one C-peptide for each insulin molecule. When a patient has newly diagnosed type 1 or type 2 diabetes, C-peptide can be used to help determine how much insulin the patient's pancreas is still producing. C-peptide measurements also can be used in conjunction with insulin and glucose levels to help diagnose the cause of documented hypoglycemia and to monitor its treatment.

PARAMETERS TESTED ON EACH LOT

Product appearance	Liquid, may turn slightly opaque during storage
Product concentration	5.0 mg/ml (+/- 10 %)
Immunoreactivity	80-120 % compared to the reference sample in an FIA test
IEF Profile	6.6 - 7.7
Purity	≥ 95 %

PARAMETERS DETERMINED DURING PRODUCT DEVELOPMENT

Subclass	IgG ₁
Association rate constant	Not Determined (N/D)
Dissociation rate constant	N/D
Affinity constant	1 x 10 ⁸ l/mol
Determination method	Radioimmunoassay (RIA)
Determination antigen	C-Peptide, Proinsulin [33-63], human C-Peptide, American Peptide Company (Cat 20-1-11, Lot 010116A1)

Legal disclaimer

Medix Biochemica products meet their specifications if transported, stored and used according to the instructions. MedixMAB is a registered trademark of Medix Biochemica and may not be used or reproduced without Medix Biochemica's written permission.

Oy Medix Biochemica Ab

Headquarters and manufacturing
Klovinpellontie 3
FI-02180 Espoo, Finland

medix@medixbiochemica.com
www.medixbiochemica.com
VAT reg. no. FI4631532

Cross-reactivities

Human proinsulin (recombinant) 18 % (Sigma, Cat P4672, Lot 5740974)

Epitope

The binding site is located within the range 4–19, with the most critical amino acids being DL (4–5).

Pair recommendations

CAPTURE ANTIBODY
9101

DETECTION ANTIBODY
9103

Please note that pair recommendations are based on results obtained by our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations are only indicative.

Product stability**TEMPERATURE, TIME**

–70 °C, 21 days

–20 °C, 21 days

+4 °C, 21 days

+25 °C, 21 days

+35 °C, 7 days

+35 °C, 21 days

+45 °C, 3 days

+45 °C, 7 days

RESULT

Failed due to reduced antigen binding

Failed due to aggregation

OK

OK

OK

Failed due to reduced antigen binding

Failed due to reduced antigen binding

Failed due to reduced antigen binding

Stability testing is performed in the product buffer to see whether different temperatures affect the antigen binding, charge or composition of the antibody. Please note that the shelf life given on the first page is based on real time stability testing at 2–8 °C in the product buffer.

Miscellaneous

–

References

–

Legal disclaimer

Medix Biochemica products meet their specifications if transported, stored and used according to the instructions. MedixMAB is a registered trademark of Medix Biochemica and may not be used or reproduced without Medix Biochemica's written permission.

Oy Medix Biochemica Ab

Headquarters and manufacturing
Klovinpellontie 3
FI-02180 Espoo, Finland

medix@medixbiochemica.com
www.medixbiochemica.com
VAT reg. no. FI4631532