

PRODUCT SPECIFICATIONS

Name	Anti-h IgG 7701 SPRN-10
Specificity	Antibody recognizes human immunoglobulin G
Description	Monoclonal mouse antibody, cultured <i>in vitro</i> under conditions free from animal-derived components
Product code	100152
Product buffer solution	37 mM citrate, 125 mM phosphate, pH 6.0, 0.9 % NaCl, 0.095 % NaN ₃ as a preservative
Shelf life and storage	36 months from manufacturing at 2-8 °C
Analyte description	Immunoglobulin G (IgG) is a monomeric immunoglobulin, built of two heavy chains and two light chains. Each IgG has two antigen binding sites. It is the most abundant immunoglobulin and is approximately equally distributed in blood and in tissue liquids, constituting 75 % of serum immunoglobulins in humans. There are four IgG subclasses (IgG1, 2, 3 and 4) in humans, named in order of their abundance in serum (IgG1 being the most abundant).

PARAMETERS TESTED ON EACH LOT

Product appearance	Liquid, may turn slightly opaque during storage
Product concentration	> 10.0 mg/ml
Immunoreactivity	80-120 % compared to the reference sample in an FIA test
IEF Profile	6.7 - 7.8
Purity	≥ 95 %

PARAMETERS DETERMINED DURING PRODUCT DEVELOPMENT

Subclass	IgG _{2a}
Association rate constant	Not Determined (N/D)
Dissociation rate constant	N/D
Affinity constant	2 x 10 ¹⁰ l/mol
Determination method	Radioimmunoassay (RIA)
Determination antigen	IgG, Scripps Laboratories (Cat I1424, Lot 484183)

Cross-reactivities

Human IgG ₁	147 % (Chemicon, Cat AG502, Lot 030BCS3)
Human IgG ₂	150 % (Chemicon, Cat AG504, Lot 212CCH)
Human IgG ₃	94 % (Chemicon, Cat AG506, Lot 030BCS4)
Human IgG ₄	2.1 % (Chemicon, Cat AG508, Lot 107BCS)
Human IgM	0.6 % (Scripps Laboratories, Cat I1124, Lot 794283)
Human IgA	0.3 % (Scripps Laboratories, Cat I2224, Lot 127283)
Goat IgG	< 0.01 % (Chemicon, Cat PP40, Lot 1250CM)
Rabbit IgG	< 0.01 % (Chemicon, Cat PP64, Lot 3080CM)
Equine IgG	< 0.01 % (Chemicon, Cat PP30, Lot 310JCI1)
Porcine IgG	< 0.01 % (Chemicon, Cat AG114, Lot 154JCI1)
Bovine IgG	< 0.01 % (Chemicon, Cat PP03, Lot 012291CHM3)
Ovine IgG	< 0.01 % (Chemicon, Cat PP44, Lot 0630CM)

Epitope

N/D

Pair recommendations

CAPTURE ANTIBODY

DETECTION ANTIBODY

-

-

Please note that pair recommendations are based on results obtained by our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations are only indicative.

Product stability

TEMPERATURE, TIME

RESULT

-70 °C, 21 days	N/D
-20 °C, 21 days	OK
+4 °C, 21 days	OK
+25 °C, 21days	N/D
+35 °C, 21 days	OK
+45 °C, 7 days	OK

Stability testing is performed in the product buffer to see whether different temperatures affect the antigen binding, charge or composition of the antibody. Please note that the shelf life given on the first page is based on real time stability testing at 2-8 °C in the product buffer.

Miscellaneous

-

References

Roivainen, M., Alfthan, G., Jousilahti, P., Kimpimäki, M., Hovi, T., and Tuomilehto, J., (1998) Enterovirus infections as a possible risk factor for myocardial infarction. *Circulation*, 98:2534-2537

Legal disclaimer

Medix Biochemica products meet their specifications if transported, stored and used according to the instructions. MedixMAB is a registered trademark of Medix Biochemica and may not be used or reproduced without Medix Biochemica's written permission.

Oy Medix Biochemica Ab

Headquarters and manufacturing
Klovinpellontie 3
FI-02180 Espoo, Finland

medix@medixbiochemica.com
www.medixbiochemica.com
VAT reg. no. FI4631532