

PRODUCT SPECIFICATIONS

Name	Anti-h Copeptin 4802 SPTN-5
Specificity	Antibody recognizes human copeptin
Description	Monoclonal mouse antibody, cultured <i>in vitro</i> under conditions free from animal-derived components
Product code	100639
Product buffer solution	50 mM Na-citrate, pH 6.0, 0.9 % NaCl, 0.095 % NaN ₃ as a preservative
Shelf life and storage	Unspecified, storage at 2-8 °C
Analyte description	Copeptin is a 39-amino acid glycopeptide, cleaved from the C-terminus of pre-provasopressin (pre-proAVP). It has been suggested as a biomarker in diagnosis and prognosis of several diseases, such as acute myocardial infarction, heart failure, hyponatremia, and sepsis.

PARAMETERS TESTED ON EACH LOT

Product appearance	Liquid, may turn slightly opaque during storage
Product concentration	5.0 mg/ml (+/- 10 %)
Immunoreactivity	80-120 % compared to the reference sample in an FIA test
IEF Profile	6.4 - 6.8
Purity	≥ 95 %

PARAMETERS DETERMINED DURING PRODUCT DEVELOPMENT

Subclass	IgG ₁
Association rate constant	Not Determined (N/D)
Dissociation rate constant	N/D
Affinity constant	N/D
Determination method	-
Determination antigen	-

Legal disclaimer

Medix Biochemica products meet their specifications if transported, stored and used according to the instructions. MedixMAB is a registered trademark of Medix Biochemica and may not be used or reproduced without Medix Biochemica's written permission.

Oy Medix Biochemica Ab

Headquarters and manufacturing
Klovinpellontie 3
FI-02180 Espoo, Finland

medix@medixbiochemica.com
www.medixbiochemica.com
VAT reg. no. FI4631532

Cross-reactivities	N/D	
Epitope	Epitope is located within the sequence ATQLDGPAGALLRLVQ in the N-terminal part of copeptin	
Pair recommendations	CAPTURE ANTIBODY 4804, 4806 4802	DETECTION ANTIBODY 4802 4804, 4806
	Please note that pair recommendations are based on results obtained by our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations are only indicative.	
Product stability	TEMPERATURE, TIME -70 °C, 21 days -20 °C, 21 days +4 °C, 21 days +35 °C, 21 days +45 °C, 7 days	RESULT OK OK OK OK OK
	Stability testing is performed in the product buffer to see whether different temperatures affect the antigen binding, charge or composition of the antibody. Please note that the shelf life given on the first page is based on real time stability testing at 2–8 °C in the product buffer.	
Miscellaneous	-	
References	-	

Legal disclaimer

Medix Biochemica products meet their specifications if transported, stored and used according to the instructions. MedixMAB is a registered trademark of Medix Biochemica and may not be used or reproduced without Medix Biochemica's written permission.

Oy Medix Biochemica Ab

Headquarters and manufacturing
Klovinpellontie 3
FI-02180 Espoo, Finland

medix@medixbiochemica.com
www.medixbiochemica.com
VAT reg. no. FI4631532