

## PRODUCT SPECIFICATIONS

<b>Name</b>	Anti-h ST2 10207 SPTN-5
<b>Specificity</b>	Antibody recognizes human soluble ST2
<b>Description</b>	Monoclonal mouse antibody, cultured <i>in vitro</i> under conditions free from animal-derived components
<b>Product code</b>	100686
<b>Product buffer solution</b>	50 mM Na-citrate, pH 6.0, 0.9 % NaCl, 0.095 % NaN <sub>3</sub> as a preservative
<b>Shelf life and storage</b>	Unspecified, storage at 2-8 °C
<b>Analyte description</b>	ST2 is an interleukin-1 family receptor expressed in the heart. ST2 protein has two isoforms: a soluble (sST2) and membrane-bound form. Patients with heart failure and elevated ST2 protein levels in the blood are at risk for heart failure progression.

## PARAMETERS TESTED ON EACH LOT

<b>Product appearance</b>	Liquid, may turn slightly opaque during storage
<b>Product concentration</b>	5.0 mg/ml (+/- 10 %)
<b>Immunoreactivity</b>	80-120 % compared to the reference sample in an FIA test
<b>IEF Profile</b>	6.2 - 6.5
<b>Purity</b>	≥ 95 %

## PARAMETERS DETERMINED DURING PRODUCT DEVELOPMENT

<b>Subclass</b>	IgG <sub>1</sub>
<b>Association rate constant</b>	Not Determined (N/D)
<b>Dissociation rate constant</b>	N/D
<b>Affinity constant</b>	N/D
<b>Determination method</b>	-
<b>Determination antigen</b>	-

### Legal disclaimer

Medix Biochemica products meet their specifications if transported, stored and used according to the instructions. MedixMAB is a registered trademark of Medix Biochemica and may not be used or reproduced without Medix Biochemica's written permission.

### Oy Medix Biochemica Ab

**Headquarters and manufacturing**  
Klovinpellontie 3  
FI-02180 Espoo, Finland

medix@medixbiochemica.com  
www.medixbiochemica.com  
VAT reg. no. FI4631532

**Cross-reactivities**

N/D

**Epitope**

Recognizes free sST2

**Pair recommendations**
**CAPTURE ANTIBODY**  
 10207  
 10201, 10202, 10203

**DETECTION ANTIBODY**  
 10201, 10202, 10203  
 10207

Please note that pair recommendations are based on results obtained by our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations are only indicative.

**Product stability**
**TEMPERATURE, TIME**
**RESULT**

-70 °C, 21 days

OK

-20 °C, 21 days

OK

+4 °C, 21 days

OK

+35 °C, 7 days

OK

+35 °C, 21 days

Reduced immunoreactivity

+45 °C, 3 days

OK

+45 °C, 7 days

Reduced immunoreactivity

Stability testing is performed in the product buffer to see whether different temperatures affect the antigen binding, charge or composition of the antibody. Please note that the shelf life given on the first page is based on real time stability testing at 2–8 °C in the product buffer.

**Miscellaneous**

Reacts with recombinant ST2 antigen, Medix Biochemica, 710020

**References**

–

**Legal disclaimer**

Medix Biochemica products meet their specifications if transported, stored and used according to the instructions. MedixMAB is a registered trademark of Medix Biochemica and may not be used or reproduced without Medix Biochemica's written permission.

**Oy Medix Biochemica Ab**

**Headquarters and manufacturing**  
 Klovipellontie 3  
 FI-02180 Espoo, Finland

medix@medixbiochemica.com  
 www.medixbiochemica.com  
 VAT reg. no. FI4631532