

PRODUCT SPECIFICATIONS

Name	Anti-h KIM-1 10103 SPTN-5
Specificity	Antibody recognizes human kidney injury molecule 1
Description	Monoclonal mouse antibody, cultured <i>in vitro</i> under conditions free from animal-derived components
Product code	100740
Product buffer solution	50 mM Na-citrate, pH 6.0, 0.9 % NaCl, 0.095 % NaN ₃ as a preservative
Shelf life and storage	Unspecified, storage at 2-8 °C
Analyte description	Kidney injury molecule 1 (KIM-1), also known as T-cell immunoglobulin mucin receptor 1 (TIM-1) or Hepatitis A virus cellular receptor 1 (HAVcr-1) is a type I transmembrane protein expressed in the renal tubular cells. KIM-1 is released after tubular injury and can be used in the diagnosis of acute kidney injury (AKI).

PARAMETERS TESTED ON EACH LOT

Product appearance	Liquid, may turn slightly opaque during storage
Product concentration	5.0 mg/ml (+/- 10 %)
Immunoreactivity	80-120 % compared to the reference sample in an FIA test
IEF Profile	6.3 - 7.2
Purity	≥ 95 %

PARAMETERS DETERMINED DURING PRODUCT DEVELOPMENT

Subclass	IgG ₁
Association rate constant	Not Determined (N/D)
Dissociation rate constant	N/D
Affinity constant	N/D
Determination method	-
Determination antigen	-

Legal disclaimer

Medix Biochemica products meet their specifications if transported, stored and used according to the instructions. MedixMAB is a registered trademark of Medix Biochemica and may not be used or reproduced without Medix Biochemica's written permission.

Oy Medix Biochemica Ab

Headquarters and manufacturing
Klovinpellontie 3
FI-02180 Espoo, Finland

medix@medixbiochemica.com
www.medixbiochemica.com
VAT reg. no. FI4631532

Cross-reactivities

No cross-reactivity with TIM-3/HAVcr-2/KIM-3 or TIM-4/TIMD-4.

Epitope

N/D

Pair recommendations

CAPTURE ANTIBODY
10103

DETECTION ANTIBODY
10101

Please note that pair recommendations are based on results obtained by our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations are only indicative.

Product stability
TEMPERATURE, TIME
RESULT

-70 °C, 21 days

OK

-20 °C, 21 days

OK

+4 °C, 21 days

OK

+35 °C, 7 days

OK

+35 °C, 21 days

Reduced immunoreactivity

+45 °C, 3 days

OK

+45 °C, 7 days

Reduced immunoreactivity

Stability testing is performed in the product buffer to see whether different temperatures affect the antigen binding, charge or composition of the antibody. Please note that the shelf life given on the first page is based on real time stability testing at 2-8 °C in the product buffer.

Miscellaneous

-

References

-

Legal disclaimer

Medix Biochemica products meet their specifications if transported, stored and used according to the instructions. MedixMAB is a registered trademark of Medix Biochemica and may not be used or reproduced without Medix Biochemica's written permission.

Oy Medix Biochemica Ab

Headquarters and manufacturing
Klovinpellontie 3
FI-02180 Espoo, Finland

medix@medixbiochemica.com
www.medixbiochemica.com
VAT reg. no. FI4631532