

Product specifications

Name	Anti-HBsAg 2507 SPTN-5
Specificity	Antibody recognizes hepatitis B virus surface antigen HBsAg preferring (ay) subtype
Description	Monoclonal mouse antibody, cultured <i>in vitro</i> under conditions free from animal-derived components.
Product code	100502
Product buffer solution	50 mM Na-citrate, pH 6.0, 0.9 % NaCl, 0.095 % NaN ₃ as a preservative
Shelf life and storage	Unspecified, storage at 2–8 °C
Subclass	IgG ₁
Analyte description	Hepatitis B is an infectious illness caused by hepatitis B virus (HBV). The acute illness causes liver inflammation, vomiting, jaundice and rarely, death. Chronic hepatitis B may eventually cause liver cirrhosis and liver cancer – a fatal disease with very poor response to current chemotherapy. The infection is preventable by vaccination. The hepatitis B surface antigen (HBsAg) is most frequently used to screen for the presence of this infection. It is the first detectable viral antigen to appear during infection. However, early in an infection, this antigen may not be present and it may be undetectable later in the infection as it is being cleared by the host.

Parameters tested on each lot

Product appearance	Liquid, may turn slightly opaque during storage
Product concentration	5.0 mg/ml (+/- 10 %)
Immunoreactivity	80–120 % compared to the reference sample in an FIA test
IEF Profile	6.3–7.0
Purity	≥ 95 %

Kinetic parameters

Association rate constant	Not Determined (N/D)
Dissociation rate constant	N/D
Affinity constant	N/D
Determination method	-
Determination antigen	-



Cross-reactivities N/D

Epitope N/D

Pair recommendations

		DETECTION			
		2505	2507	2508	2510
CAPTURE	2505	-	+	-	+
	2507	-	+	+	+
	2508	+	+	+	+
	2510	+	+	+	+

Antibody 2507 prefers HBsAg (ay) subtype.

Please note that pair recommendations are based on results obtained by our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations are only indicative.

Platforms tested FIA, LF

Antigens tested N/D

TEMPERATURE, TIME	RESULT
-70 °C, 21 days	OK
-20 °C, 21 days	OK
+4 °C, 21 days	OK
+35 °C, 7 days	OK
+35 °C, 21 days	Charge alterations
+45 °C, 3 days	Charge alterations
+45 °C, 7 days	Charge alterations

Stability testing is performed in the product buffer to see whether different temperatures affect the antigen binding, charge or composition of the antibody. Please note that the shelf life given on the first page is based on real time stability testing at 2–8 °C in the product buffer.

Miscellaneous -

References -

